

Enzymax, LLC
870 Corporate Drive, suite 201
Lexington, KY 40503

Product Information

Product Name: Human DCP2
Catalog #: 86
Size: 10 µg

Order: info@enzymax.net
Tel: 859-219-8482
Fax: 859-219-0653
Web: www.enzymax.net

Product: human DCP2 decapping enzyme

Product Description:

Human DCP2 is a decapping enzyme of mRNA, which is a critical step in eukaryotic mRNA turnover. This enzyme specifically hydrolyze methylated capped RNA (m7 GTP caps) to release m7GDP and 5' monophosphorylated RNA.

Source: Recombinant human DCP2 was expressed in and purified from *E. coli*.

Molecular Weight:

Full length hDCP2 (~69 kDa)

Decapping Reaction Buffers: 1X


10 mM Tris-HCl at pH 7.5
100 mM NaCl
2 mM MgCl₂
1 mM DTT
1 mM MnCl₂

10x decapping Reaction Buffer: Included

NOTE: The following two components are option based on your applications:
40 units/mL (recombinant RNase inhibitor)
0.1 mM Spermine

Presentation: Purified hDCP2 is supplied at a concentration range from 0.1-0.5 mg/ml in a buffer containing 50 mM Tris-HCl (pH 8.0), 500 mM NaCl and 10% glycerol.

SDS-PAGE gel picture:



Purified human DCP2. DCP2 (300 ng) was analyzed by electrophoresis on a 15% SDS-polyacrylamide gel and visualized by staining with Coomassie blue. Protein size markers are indicated on the left.

Suggested condition for used in 5'RACE:
It may not be optimized for other applications.

Reaction temperature/time: 37°C, 30-120 min
Reaction volume: 20-50 µl
RNA substrates: 100ng to 1ug
DeCapping enzymes: 50-500ng (~1-10 pmol)

Contaminating Activity Assay: It is free of detectable RNase activity when incubation of 2ug of SP6 transcript with 50ng of enzyme for 1 hour at 37°C. It is also free of detectable exo and endo nuclease activities.

Storage: Stable for 12 months at -80°C from date of shipment. Please aliquot to avoid repeated freezing and thawing.